



| Question | Responses |
|---|---|
| Demographics | • Age |
| | Gender |
| How many suspected non- melanoma skin cancer lesions are being excised? | • 1, 2, 3, 4, 5 o If 2 or more is selected, then all questions below will be repeated for the 2 nd , 3 rd lesions etc |
| Pre-op diagnosis (from Plastic Surgery) | BCCSCCUnknown or biopsy proven |
| Is the lesion ulcerated? | Yes, No, Unknown |
| Is this area within 10mm of a previous treatment scar? | Head and neck Scalp Periocular, temple, forehead or eyebrow Cheek or chin Ear (or within 2cm) Nose (or within 1cm) or lips Neck Trunk Limb Hand or foot Genitalia, perineum or perianal area Unknown Yes, No, Unknown |
| Type of excision | Date of surgery Primary Re-excision Unknown |
| Planned clinical margin from op | Peripheral (mm or unknown) Poor (mm tissue layer or unknown) |
| Type of immediate reconstruction | Deep (mm, tissue layer or unknown) Left open (or delayed reconstruction) Direct closure Skin graft Full thickness skin graft Split thickness skin graft Unknown Flap Unknown |
| Did the patient receive prophylactic antibiotics? | At induction Yes, No, Unknown Post-op |

| | Yes, No, Unknown |
|-----------------------------------|---|
| Histological diagnosis | BCC Subtypes – can select more than one if mixed Nodular Infiltrative Superficial Micronodular Morpheaform Sclerosing Adenoid Fibroepithliomatous Other Unknown SCC Actinic keratosis Bowen's Disease |
| | Melanoma Benign – therefore skip next 2 questions Other |
| Largest tumour diameter | Size in mm (or unknown) |
| Margins (from histology report) | Peripheral (mm or involved) Deep margin (mm or involved) |
| Complications - Definitions above | None Bleeding Infection Graft/flap failure for other reasons Other Unknown / patient not followed up |
| Further treatment | Listed for re-excision Yes, No, Unknown Referral for radiotherapy Yes, No, Unknown Discharged from Plastic Surgery? |